

In vitro Variation in antibacterial activity plant extracts on *Glaucium elegans* and Saffron (*Crocus sativus* .L) Onios

Ehsan Heidari Soureshjan^{1,*}, Mina Heidari^{1,2}

¹ Department of Biology, Young Researchers and Elite Club, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran;

² Department of Biology, Shahrekord University, Shahrekord, Iran.

* Corresponding author. Tel: +98-38-13361001, +98-9139802276; E-mail: ehsanheidari2012@yahoo.com, ehsan.heidari@iaus.ac.ir

Abstract

The increase in antibiotic resistance has resulted in a decreasing number of fully active antimicrobial agents available to treat infections by multi-drug resistant (MDR) bacteria. This has necessitated a search for new antimicrobial agents. Though these plants are used for various purposes, but there are plants that the medicinal properties of the plants were not investigated. The major aim of this study was to determine the antimicrobial activity of the extracts of *Glaucium elegans* and Saffron (*Crocus sativus* .L) Onios plant species against *Escherichia coli*, *Staphylococcus aureus*, *Salmonella enteritidis*, *Bacillus anthracis* and *Proteus*. The extracts of medicinal plants were prepared using Soxhlet apparatus for methanol extract. The antibacterial activities by agar disc diffusion and serial dilution assays; the inhibitory zones were recorded in millimeters. The methanol extract of *Glaucium elegans* was found to have a significant antibacterial efficiency ($p \leq 0.05$) as compared to the methanol extract of Onios plant. The methanol extract of *Glaucium elegans* showed high antibacterial potential (23 mm) against *Proteus* at concentration of 1500 μ g/disc. On the other hand showed highest antibacterial activities (22 mm) against *Staphylococcus aureus* at concentration of 1500 μ g/disc. These finding pinpoint the efficiency of these extracts to inhibit microbial growth. The bactericidal activity described here represents an added safety value for *Glaucium elegans* as a means of eliminating a possible antimicrobial during storage.

Keywords: *Glaucium elegans*, Saffron Onios, *Escherichia coli*, *Staphylococcus aureus*, *Salmonella enteritidis*, *Bacillus anthracis*, *Proteus*.

1. Introduction

For over thousands of years now, natural plants have been seen as a valuable source of medicinal agents with proven potential of treating infectious

diseases and with lesser side effects compared to the synthetic drug agents (1). Numerous studies have shown that aromatic and medicinal plants are sources of diverse nutrient and non nutrient molecules, many of which display antioxidant and antimicrobial properties which can protect the human body against both cellular oxidation reactions and pathogens. Thus it is important to characterize different types of medicinal plants for their antioxidant and antimicrobial potential (2-4). The world health organisation estimates that 80% of the people living in developing countries almost exclusively use traditional medicine. Medicinal plants form the principle component of traditional medicine (5). Aromatic and medicinal plants are known to produce certain bioactive molecules which react with other organisms in the environment, inhibiting bacterial or fungal growth (antimicrobial activity) (6). The genus *Glaucium* Mill. Belongs to *Papaveraceae* family and consists of 25 species, annuals, biennials and perennials. They grow in middle and South West Asia and Europe (7). This genus consists of herbaceous plants. Cullen (1966) in Flora Iranica area recognized 11 species, 10 of which occurring in Iran, one including two subspecies. Mobayen (1984) described two new species, namely *G. elegantissimum* Mobayen and *G. mathiolifolium* Mobayen. Later on Mobayen (1985) in Flora of Iran, vascular plants apparently recognized 14 species in Iran, several of them including subspecific taxa (8). Saffron is the most expensive spice in the world and consists of the dried stigmas of *Crocus sativus* L. By far the most important producer is Iran, followed by Greece, Morocco, India, Spain and Italy. Saffron is greatly appreciated for its abilities of colouring and flavouring, and for its aromatic strength. These properties are basically related to its contents of picrocrocin, safranal and crocins (9). Picrocrocin is the glycoside precursor of safranal (2, 6, 6-trimethyl-1, 3-cyclohexadiene-1-carboxaldehyde), which is in turn the most abundant of the volatile compounds responsible for the aroma of this spice (10).

Spices have been used since ancient times to hide spoilage in foods. Nonetheless, as with many

other agricultural products, spices can suffer microbial contamination and their additions to ready-to eat foods can result in the proliferation of bacterial pathogens (11,12). The wide use of antibiotics in the treatment of bacterial infections has led to the emergence and spread of resistant strains (13). The antimicrobial compounds from plants may inhibit bacteria by a different mechanism than the presently used antibiotics and may have clinical value in treatment of resistant microbial strains. Because of the side effects and bacteria resistance against the antibiotics, the scientist developed new drugs from natural sources such as plants, which have been extensively used as alternative treatment for disease (14, 15). The purpose of the present study was to investigate antimicrobial properties of *Glaucium elegans* and Saffron Onion. In this paper we report the results of such studies in order to orient future investigations towards the finding of new, potent and safe antimicrobial compounds.

2. Materials and Methods

2.1 Plant materials

Glaucium elegans and Saffron Onion were collected from different natural habitats of the country. The Onion of Saffron and Leaves and twigs of *Glaucium elegans* separated and dried in dark at room temperature. In order to extract content of 12 g fresh sample prepared with distilled water and poured into an Erlenmeyer flask and heated for 30 min at 100 ° C was boiled. The extracts were filtered using Whitman Strainer paper and then in Dry Heat of 55 ° C for 24 h were exposed to the extract powder is produced. One gram of the powder, each of which was dissolved in 10 ml of Deionized water. Different concentration (250, 750 and 1500 µg/disc) and the extracts were prepared calculated.

2.2 Bacterial strains and culture media

Five pathogenic bacteria including five strains of *Staphylococcus aureus*, *Escherichia coli*, *Salmonella enteritidis*, *Bacillus anthracis* and *Proteus* were used for the bioassay study. The pure strain was identified and obtained from Microbiology and Biotechnology Laboratory, Department of Islam Islamic Azad University, Shahrekord Branch, Shahrekord, Iran. All cultures were incubated at 37 °C in aerobic conditions. The test organisms were transferred to the agar slants from the supplied pure cultures with the help of an inoculating loop in an aseptic condition. The inoculated slants were then incubated at 37.5 °C for 24 hrs to ensure the growth of test organisms. These fresh cultures were used for the sensitivity test.

2.3 Preparation of discs

Three types of discs were prepared for antibacterial screening. Sterilized (BBL, USA) filter paper discs (5 mm in diameter) were taken in blank Petri dishes. Sample solution of desired concentration (10µl/ disc) was applied on the discs with the help of a micropipette in an aseptic condition. These were used to compare the antibacterial activity of test material. In our investigation, Gentamicin (1.25 µg/disc) was used as standard disc. These were prepared by using identical filter paper (5 mm diameter) and same volume of residual solvent in the same condition. These were used as negative control to ensure that the residual solvent and the filter paper themselves was not active.

2.4 Placement of the discs and incubation

The dried crude extract separate discs and standard discs were placed on the solidified agar plates seeded with the test organisms. The plates were kept in a refrigerator at 4 °C for 24 hrs in order to provide sufficient time to diffuse the antibiotics into the medium. Then the plates were incubated at 37.5 ° C for 24 hrs in an incubator.

2.5 Measurement of the zones of inhibition

After incubation, the antibacterial activities of the test samples were determined by measuring the diameter of inhibitory zones (mm) with a transparent scale.

2.6 Statistical analysis

The data were analyzed using Analysis of Variance (ANOVA) test. A one-way ANOVA was performed on all results. Probabilities of less than 0.05 were considered statistically significant. Differences between means were determined using Duncan compromise test. The software SPSS (Ver. 17) was used to conduct all the statistical analysis.

3. Results and Discussion

The tests for antibacterial effect establish of vegetal extracts revealed that most of the extracts are representing an important source of substances with antimicrobial activity. The results of different studies provide evidence that some medicinal plants might indeed be potential sources of new antibacterial agents (16). The antimicrobial activity of extract having different concentrations (250µg/disc, 750µg/disc and 1500µg/disc), was tested against five pathogenic bacteria. Gentamicin disc (1.25µg/disc) was used for comparing the antibacterial activity. Tables 1 represent the inhibition zone, found by the activity of extracts on microorganisms tested and compared with Gentamicin. At 250 µg/disc dosage extract of *Glaucium elegans* showed 1, 2 and 1 mm zone of inhibition against *E. coli*, *Proteus* and *B. anthracis* respectively. However no antimicrobial activity was observed against *S. aureus* and *S. enteritidis* the same concentration.


Table 1. Inhibition areas diameter of the tested plant essential extracts on microorganisms tested and compared with Gentamicin.

Name Vegetal	Concentration (µg/disc)	Test bacteria				
		E. coli	Proteus	S. aureus	S. enteritidis	B. anthracis
Glaucium elegans	250	1 ±0.3	2±0.2	-	-	1±0.1
	750	10±0.4	18±0.3	12±0.55	9±0.1	17±0.2
	1500	13±0.4	23±0.65	22±0.45	13±01	21±0.8
Saffron Onion	250	-	-	-	-	-
	750	-	-	-	-	-
	1500	-	-	-	-	-
Gentamcin	1.25	27±0.50	17±0.60	26±0.40	25±0.48	28±0.68

At 750µg/disc Concentration of *G. elegans*, the zone of inhibition against the same bacteria was 10, 18, 12, 9 and 17mm respectively against *E. coli*, *Proteus*, *S. aureus*, *S. enteritidis* and *B. anthracis* respectively. In case of highest dose of 1500 µg/disc dosage extract of *G. elegans*, the zones of inhibition were found to be 13, 23, 22, 13 and 21mm respectively (Figure 1). At all concentrations dosage extract of Saffron Onion no antimicrobial activity was observed against bacterias. But the standard antibiotic disc Gentamycin (1.25 µg/disc) showed the zone of inhibition against five bacteria such as *E. coli*, *Proteus*, *S. aureus*, *S. enteritidis* and *B. anthracis*. The zones of inhibitions were 27, 19, 26, 25 and 28mm respectively. Tested essential extracts most antimicrobial effect of the *Proteus* strains including the strains with resistance to antibiotics. It can be concluded that *G. elegans* has a wide range of antibacterial activity as medicines. Also in research on extract of *S. mirzayanii* demonstrated antimicrobial effect against *S. aureus* and *E. coli* (17). In other study assay, the methanol extract of red variety was found to have a significant antibacterial effect against *S. aureus*, *E. coli* (18). In summary, the results obtained in this work indicate that *G. elegans* extracts could be an important source of antibiotics compounds potentially active against bacterias. The *G. elegans* extract assayed has been demonstrated to be useful against *E. coli*, *Proteus*, *S. aureus*, *S. enteritidis* and *B. anthracis* strains with antibiotic resistances.

Acknowledgment

We would like to thank head and deputy of research of Islamic Azad University of Shahrekord branch in Iran and Young Researchers and Elites Club for their sincere support.

**Figure 1.** Antimicrobial activity extract of Glaucium elegans.

References

- [1] Nagham A. (2011) Study Effect of Essential Oils on Resistant Bacteria. *J. Kerbala Univ. Sci.* **9**: 367-369.
- [2] Mothana RAA, Lindequist U. (2004) Antimicrobial activity of some medicinal plants of the island Soqotra. *J. of Ethnopharmacology* **96**: 177-181.
- [3] Bajpai M, Pande A, Tewari SK and Prakash D. (2005) Phenolic contents and antioxidant activity of some food and medicinal plants. *International Journal of Food Sciences and Nutrition*, **56**: 287-291.
- [4] Wojdyło A, Oszmian´ski J, Czemerys R. (2007) *Antioxidant activity and phenolic compounds in 32 selected herbs*. Elsevier. **105**: 940–949.
- [5] Farnsworth NR. (1994) Ethnopharmacology and drug development. In: Prance, G.T. (Ed.), *Ethnobotany and the Search for New Drugs*. Wiley, Chichester (Ciba Foundation Symposium. **185**: 42–59.

- [6] Sezai E, Memnune S, Hilal Y, Neva Gr, Buent Cn, Zeynep E. (2009) Total Phenolic Content, antioxidant and antimicrobial activities of some medicinal plants. *Pak. J. Pharm. Sci.* **22**: 102-106.
- [7] Burnie G, Forrester S, Greig D, Guest S, Harmony M, Hobly S, Jackson G. (2004) *Botanica*. -Publisher Goldon Cheers, Australia.
- [8] Gran A, Sharifnia F. (2008) Micro-Macromorphological Studies of the Genus *Glaucium* (Papaveraceae) In Iran. *Iran. Journal. Bot.* **14**: 23-38.
- [9] Rafael R, Concepción P, Alicia de M, Olga A, Leonor N, José Luis N. (2011) Bactericidal effect of saffron (*Crocus sativus* L.) on *Salmonella enteric* during storage. *Food Control.* **22**: 638-642.
- [10] Maggi L, Carmona M, del Campo CP, Kanakis CD, Anastasaki EG, Tarantilis PA, et al. (2009) Worldwide market screening of saffron volatile composition. *Journal of the Science of Food and Agriculture.* **89**: 1950-1954.
- [11] Sagoo SK, Little CL, Greenwood M, Mithani V, Grant KA, McLauchlin J, et al. (2009) Assessment of the microbiological safety of dried spices and herbs from production and retail premises in the United Kingdom. *Food Microbiology* **26**: 39-43.
- [12] Vij V, Ailes E, Wolyniak C, Angulo FJ, Klontz, KC. (2006). Recalls of spices due to bacterial contamination monitored by the U.S. Food and Drug Administration: the predominance of *Salmonellae*. *Journal of Food Protection* **69**: 233-237.
- [13] Patterson JE. (2000) New gram-positive agents in L. Ran, 2005. The synergistic activity of antibiotics nosocomial infection. *Curr Opin Dis.* **13**: 593-598.
- [14] Sumitra C, Yogesh B. (2010) Extraction of Active Compounds of Some Medical Plants. *African journal of Biotechnology.* **9**: 3210-3217.
- [15] Manoj M, Kailas C, Balaji V, Sajid N. (2010) Effect of Plants Extracts. *International Journal of Pharm Tech Res.* **2**: 899-901.
- [16] Kone WM, Atindehou KK, Terreaux C, Hostettmann K, Traore D, Dosso M. (2004). Traditional medicine in North Coted'Ivoire: screening of 50 medicinal plants for antibacterial activity. *Journal of Ethnopharmacology* **93**: 43-49.
- [17] Moshafi MH, Mehrbani M, Zvalhsb H. (2004) Effects Antimicrobial *Salvia mirzayanii* and *Salvia atropatana* on six strains of gram positive and negative bacteria. *JKUMS.* **11**: 109-118.
- [18] Krishna BK , Faruk MO, Amin MZ, Shaha RK. (2012) Antibacterial activity of two varieties of water chestnuts (TRAPA SP). *J.bio-sci.* **20**: 115-123..